



YOUR CROP PRODUCTION PARTNERS™

# CONFINE™

For the post - harvest treatment of russet – skinned potatoes and potatoes intended for processing for the suppression of late blight and pink rot storage infection.

<b>GROUP</b>	<b>33</b>	<b>FUNGICIDE</b>
--------------	-----------	------------------

LIQUID  
AGRICULTURAL

REGISTRATION NO. 29100 PEST CONTROL PRODUCTS ACT

GUARANTEE: Mono- and di-potassium salts of Phosphorous Acid 45.8%

READ THE LABEL BEFORE USING  
KEEP OUT OF REACH OF CHILDREN

NET CONTENTS: \_\_\_\_\_

Winfield Solutions LLC  
P.O. Box 64589  
St. Paul, MN 55164-0589

The Agronomy Company of Canada  
17554 Plover Mills Road  
Thorndale, ON N0M 2P0  
519-461-9057

**PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN**

Avoid breathing vapors or spray mist. May cause minimal eye irritation. Avoid contact with skin, eyes or clothing. Applicators and other handlers must wear protective eyewear, long pants and long-sleeved shirt, waterproof gloves, shoes plus socks. Remove contaminated clothing and wash clothing before reuse.

**ENVIRONMENTAL HAZARDS:** DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of waste.

**FIRST AID:** If swallowed - Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing - Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled - Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes - Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**TOXICOLOGICAL INFORMATION:** Treat symptomatically.

**DISPOSAL:** Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

**NOTICE TO USER:** This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this

product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

**RESISTANCE MANAGEMENT RECOMMENDATIONS:** For resistance management, please note that CONFINE Fungicide contains a Group 33 fungicide. Any fungal population may contain individuals naturally resistant to CONFINE Fungicide and other Group 33 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same sites. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance management strategies should be followed.

To delay fungicide resistance:

- Where possible, rotate the use of CONFINE Fungicide or other Group 33 fungicides with different groups that control the same pathogens.
- Avoid application of consecutive sprays of CONFINE or other fungicides in the same group in a season.
- Fungicide use should be based on an IPM program that includes scouting, historical information related to pesticide use and crop rotation and considers cultural, biological and other chemical control practices.
- Monitor treated fungal populations for sign of resistance development. If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product, and switch to another fungicide with a different target site of action, if available.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and disease problems in your area.

For further information or to report suspected resistance, contact company representatives at 519-461-9057 or at [www.agromartgroup.com](http://www.agromartgroup.com).

**DIRECTIONS FOR USE:** Potatoes (Post-harvest treatment)

For the suppression of Late Blight (*Phytophthora infestans*) and Pink Rot (*Phytophthora erythroseptica*) on harvested potato tubers. Dilute Confine at a 1:4.3 ratio with water. Apply 2 litres of this solution as a spray to 1000 kg of potatoes prior to storage. Ensure complete and even coverage. Use only on russet-skinned varieties, or potatoes intended for processing.